arXiv:1601.02989v1 [math.CV] 12 Jan 2016

THE BERGMAN KERNEL: EXPLICIT FORMULAS, DEFLATION, LU QI-KENG PROBLEM AND JACOBI POLYNOMIALS

Tomasz Beberok

In this paper we investigate the Bergman kernel function for intersection of two complex ellipsoids

$$\{(z,w_1,w_2)\in\mathbb{C}^{n+2}:|z_1|^2+\cdots+|z_n|^2+|w_1|^q<1,\quad |z_1|^2+\cdots+|z_n|^2+|w_2|^r<1\}.$$

Keyword: Lu Qi-Keng problem, Bergman kernel, Routh-Hurwitz theorem, Jacobi polynomials

AMS Subject Classifications: 32A25; 33D70

1 Introduction

Let D be a bounded domain in $\mathbb{C}^n$. The Bergman space $L^2_a(D)$ is the space of all square integrable holomorphic functions on D. Then the Bergman kernel $K_D(z,w)$ is defined [2] by

$$K_D(z,w)=\sum_{j=0}^{\infty}\Phi_j(z)\overline{\Phi_j(w)},\quad (z,w)\in D\times D,$$

where $\{\Phi_j(\cdot)\colon j=0,1,2,...\}$ is a complete orthonormal basis for $L^2_a(D)$. It is defined for arbitrary domains, but it is hard to obtain concrete representations for the Bergman kernel except for special cases like a Hermitian ball or polydisk. Refer to [6] for more on this topic.

In 2015 [1] the author of this paper computed the Bergman kernel for $D_1^{q,r}=\{z=(z_1,w_1,w_2)\in\mathbb{C}^3\colon |z_1|^2+|w_1|^q<1,\quad |z_1|^2+|w_2|^r<1\}$ explicitly. The goal of this paper is to extend the result in [1] to higher dimensional case, namely to $D_n^{q,r}=\{(z,w_1,w_2)\in\mathbb{C}^{n+2}:|z_1|^2+\cdots+|z_n|^2+|w_1|^q<1,\quad |z_1|^2+\cdots+|z_n|^2+|w_2|^r<1\}$.

This paper will be organized as follows. In Section 2, we will compute explicit formula of the Bergman kernel for $D_n^{q,r}$ and we show deflation identity between domains $D_n^{q,r}$ and $D_{1/n}^{q,r}=\{z=(z_1,w_1,w_2)\in\mathbb{C}^3\colon |z_1|^{2/n}+|w_1|^q<1,\quad |z_1|^{2/n}+|w_2|^r<1\}$. In Section 3, we show some relation between Bergman kernel for $D_n^{2,2}$ and Jacobi polynomials. In Section 4, we investigate the Lu Qi-Keng problem for $D_n^{2,2}$. In the final section, we consider the Bergman kernel for $\Omega_n^r:=\{(z,w)\in\mathbb{C}\times\mathbb{C}^n\colon |z|^2+|w_1|^r<1,\ldots,|z|^2+|w_n|^r<1\}$.

2 Bergman Kernel

Let $\zeta = (z, w_1, w_2) \in \mathbb{C}^{n+2}$. Put $\Phi_\kappa(\zeta) = \zeta^\kappa = z^\alpha w_1^{\gamma_1} w_2^{\gamma_2}$ for each multi-index $\kappa = (\alpha, \gamma_1, \gamma_2)$. Since $D_n^{q,r}$ is complete Reinhardt domain, the collection of $\{\Phi_\kappa\}$ such that each $\alpha_i \geq 0$ and $\gamma_j \geq 0$ is a complete orthogonal set for $L^2(D_n^{q,r})$.

Proposition 2.1 *Let $\alpha_i \in \mathbb{Z}_+$ for $i = 1, \ldots, n$ and $\gamma_1 \geq 0$, $\gamma_2 \geq 0$. Then, we have*

$$\|z^\alpha w_1^{\gamma_1} w_2^{\gamma_2}\|^2_{L^2(D_n^{q,r})} = \frac{\pi^{n+2}\Gamma(\frac{2\gamma_1+2}{q} + \frac{2\gamma_2+2}{r} + 1)\prod_{i=1}^n \Gamma(\alpha_i+1)}{(\gamma_2+1)(\gamma_2+1)\Gamma(\frac{2\gamma_1+2}{q} + \frac{2\gamma_2+2}{r} + |\alpha| + n + 1)},$$

where $|\alpha| = \alpha_1 + \cdots + \alpha_n$.

Proof.

$$\|z^\alpha w_1^{\gamma_2} w_2^{\gamma_3}\|^2_{L^2(D_n^{q,r})} = \int\limits_{D_n^{q,r}} |z|^{2\alpha} |w_1|^{2\gamma_2} |w_2|^{2\gamma_3}\, dV(z) dV(w)$$

We introduce polar coordinate in each variable by putting $z = re^{i\theta}$, $w_1 = s_1 e^{i\lambda_1}$, $w_2 = s_2 e^{i\lambda_2}$. After doing so, and integrating out the angular variables we have

$$(2\pi)^{n+2} \int_{Re(D_n^{q,r})} r^{2\alpha+1} s_1^{2\gamma_1+1} s_2^{2\gamma_2+1}\, dV(r) dV(s),$$

where $Re(D_n^{q,r}) = \big\{(r,s) \in \mathbb{R}^n_+ \times \mathbb{R}^2_+ : \|r\|^2 + s_1^q < 1, \quad \|r\|^2 + s_2^r < 1\big\}$. Next we use spherical coordinates in the r variable to obtain

$$(2\pi)^{n+2} \int_{S * Re(D_n^{q,r})} \int_{\mathbf{S}_+^{n_1-1}} \rho^{2|\alpha|+2n-1} \omega^{2\alpha+1} s_1^{2\gamma_1+1} s_2^{2\gamma_2+1}\, d\sigma(\omega) dV(s) d\rho,$$

where $S * Re(D_n^{q,r}) = \big\{(\rho, s) \in \mathbb{R}_+ \times \mathbb{R}^2_+ : \rho^2 + s_1^q < 1, \quad \rho^2 + s_2^r < 1\big\}$. Using (6) form Lemma 1 in [4] we obtain

$$(2\pi)^{n+2} \frac{\beta(\alpha+1)}{2^{n-1}} \int_0^1 \int_0^{(1-\rho^2)^{1/q}} \int_0^{(1-\rho^2)^{1/r}} \rho^{2|\alpha|+2n-1} s_1^{2\gamma_1+1} s_2^{2\gamma_2+1}\, ds_1 ds_2 d\rho,$$

where $\beta(\alpha) = \frac{\prod_{i=1}^n \Gamma(\alpha_i)}{\Gamma(\alpha_1+\cdots+\alpha_n)}$. Integrating out of s_1 and s_2 variables, we have

$$\frac{(2\pi)^{n+2}\beta(\alpha+1)}{2^{n+1}(\gamma_1+1)(\gamma_2+1)} \int_0^1 \rho^{2|\alpha|+2n-1} (1-\rho^2)^{\frac{2\gamma_1+2}{q} + \frac{2\gamma_2+2}{r}}\, d\rho$$

After a little calculation using well known fact

$$\int_0^1 x^a (1-x^p)^b\, dx = \frac{\Gamma((a+1)/p)\Gamma(b+1)}{p\Gamma((a+1)/p + b + 1)},$$

we obtain desired result.

Now we discuss the Bergman kernel for $D_n^{q,r}$.

Theorem 2.1 *The Bergman kernel for $D_n^{q,r}$ is given by*

$$K_{D_n^{q,r}}((z,w_1,w_2),(\eta,\xi_1,\xi_2)) = \frac{L_n^{q,r}(a,b)}{\pi^{n+2}(1-z_1\overline{\eta}_1-\cdots-z_n\overline{\eta}_n)^{\frac{2}{q}+\frac{2}{r}+1}},$$

where

$$a = \frac{w_1\overline{\xi}_1}{(1-z_1\overline{\eta}_1-\cdots-z_n\overline{\eta}_n)^{\frac{2}{q}}}, \quad b = \frac{w_2\overline{\xi}_2}{(1-z_1\overline{\eta}_1-\cdots-z_n\overline{\eta}_n)^{\frac{2}{r}}}$$

$$L_{n+1}^{q,r}(x,y) = nL_n^{q,r}(x,y) + \frac{2}{q}\frac{\partial}{\partial x}xL_n^{q,r}(x,y) + \frac{2}{r}\frac{\partial}{\partial y}yL_n^{q,r}(x,y) \text{ and}$$

$$L_1^{q,r}(x,y) = \frac{2(q(1-x)(y+1)+r(x+1)(1-y))}{qr(1-x)^3(1-y)^3}$$

Proof. By Proposition 2.1, we have

$$K_{D_n^{q,r}}((z,w_1,w_2),(\eta,\xi_1,\xi_2))$$
$$= \frac{1}{\pi^{n+2}}\sum_{\alpha=0}^{\infty}\sum_{\gamma_1,\gamma_2=0}^{\infty}\frac{(\gamma_1+1)(\gamma_2+1)\Gamma(\frac{2\gamma_1+2}{q}+\frac{2\gamma_2+2}{r}+|\alpha|+n+1)}{\Gamma(\frac{2\gamma_1+2}{q}+\frac{2\gamma_2+2}{r}+1)\prod_{i=1}^{n}\Gamma(\alpha_i+1)}\mu^{\alpha}\nu_1^{\gamma_1}\nu_2^{\gamma_2},$$

where $\mu^{\alpha} = (z_1\overline{\eta}_1)^{\alpha_1}\cdot\cdots\cdot(z_n\overline{\eta}_n)^{\alpha_n}$ and $\nu_1 = w_1\overline{\xi}_1$, $\nu_2 = w_2\overline{\xi}_2$. Sum out of each α_i, we have

$$\frac{1}{\pi^{n+2}(1-\tau)^{\frac{2}{q}+\frac{2}{r}+1}}\sum_{\gamma_1,\gamma_2=0}^{\infty}\frac{(\gamma_1+1)(\gamma_2+1)\Gamma(\frac{2\gamma_1+2}{q}+\frac{2\gamma_2+2}{r}+n+1)}{\Gamma(\frac{2\gamma_1+2}{q}+\frac{2\gamma_2+2}{r}+1)(1-\tau)^{\frac{2\gamma_1}{q}+\frac{2\gamma_2}{r}}}\nu_1^{\gamma_1}\nu_2^{\gamma_2},$$

where $\tau = z_1\overline{\eta}_1+\cdots+z_n\overline{\eta}_n$. Now we will consider sequence of functions $L_n^{q,r}$ defined as follows

$$L_n^{q,r}(x,y) = \sum_{\gamma_1,\gamma_2=0}^{\infty}\frac{(\gamma_1+1)(\gamma_2+1)\Gamma(\frac{2\gamma_1+2}{q}+\frac{2\gamma_2+2}{r}+n+1)}{\Gamma(\frac{2\gamma_1+2}{q}+\frac{2\gamma_2+2}{r}+1)}x^{\gamma_1}y^{\gamma_2}$$

Using the identity $\Gamma(t+1) = t\Gamma(t)$ we easily obtain recursion formula

$$L_{n+1}^{q,r}(x,y) = nL_n^{q,r}(x,y) + \frac{2}{q}\frac{\partial}{\partial x}xL_n^{q,r}(x,y) + \frac{2}{r}\frac{\partial}{\partial y}yL_n^{q,r}(x,y)$$

Moreover

$$L_1^{q,r}(x,y) = \sum_{\gamma_1,\gamma_2=0}^{\infty}(\gamma_1+1)(\gamma_2+1)\left(\frac{2\gamma_1+2}{q}+\frac{2\gamma_2+2}{r}+1\right)x^{\gamma_1}y^{\gamma_2}$$

Hence

$$L_1^{q,r}(x,y) = \frac{2(q(1-x)(y+1)+r(x+1)(1-y))}{qr(1-x)^3(1-y)^3}$$

which completes the proof.

2.1 Deflation

Now we will consider following domains

$$D^{q,r}_{1/n} = \{(z, w_1, w_2) \in \mathbb{C}^3 \colon |z|^{2/n} + |w_1|^q < 1, \quad |z|^{2/n} + |w_2|^r < 1\}$$

Similarly as in Sect. 2, we have

Proposition 2.2 *Let $\alpha \geq 0$, $\gamma_1 \geq 0$ and $\gamma_2 \geq 0$. Then, we have*

$$\|z^\alpha w_1^{\gamma_1} w_2^{\gamma_2}\|^2_{L^2(D^{q,r}_{1/n})} = \frac{n\pi^3 \Gamma(\frac{2\gamma_1+2}{q} + \frac{2\gamma_2+2}{r} + 1)\Gamma(n\alpha + n)}{(\gamma_2+1)(\gamma_2+1)\Gamma(\frac{2\gamma_1+2}{q} + \frac{2\gamma_2+2}{r} + n\alpha + n + 1)}.$$

Therefore

$$\begin{aligned} &K_{D^{q,r}_{1/n}}((0, w_1, w_2), (0, \xi_1, \xi_2)) \\ &= \sum_{\gamma_1,\gamma_2=0}^{\infty} \frac{(\gamma_1+1)(\gamma_2+1)\Gamma(\frac{2\gamma_1+2}{q} + \frac{2\gamma_2+2}{r} + n + 1)}{\pi^3 n! \Gamma(\frac{2\gamma_1+2}{q} + \frac{2\gamma_2+2}{r} + 1)} (w_1\overline{\xi}_1)^{\gamma_1} (w_2\overline{\xi}_2)^{\gamma_2} \end{aligned}$$

Hance

$$K_{D^{q,r}_{1/n}}((0, w_1, w_2), (0, \xi_1, \xi_2)) = \frac{1}{\pi^3 n!} L^{q,r}_n(w_1\overline{\xi}_1, w_2\overline{\xi}_2)$$

Comparing formulas for $K_{D^{q,r}_{1/n}}$ and $K_{D^{q,r}_n}$, we obtain following deflation identity

Proposition 2.3 *For every $n \in \mathbb{N}$ and every positive numbers q and r, we have*

$$\frac{n!}{\pi^{n-1}} K_{D^{q,r}_{1/n}}((0, w_1, w_2), (0, \xi_1, \xi_2)) = K_{D^{q,r}_n}((0, \ldots, 0, w_1, w_2), (0, \ldots, 0, \xi_1, \xi_2))$$

Note that, we have some kind of deflation result similar to that obtained in [3].

3 Some representations of Bergman kernel for $D^{2,2}_n$

Jacobi polynomials are a class of classical orthogonal polynomials. They are orthogonal with respect to the weight $(1-x)^k(1+x)^l$ on the interval $[-1, 1]$. For k, $l > -1$ the Jacobi polynomials are given by the formula

$$P^{(k,l)}_d(z) = \frac{(-1)^d}{2^d d!}(1-z)^{-k}(1+z)^{-l}\frac{\partial}{\partial z^d}\left\{(1-z)^k(1+z)^l(1-z^2)^d\right\}.$$

The Jacobi polynomials are defined via the hypergeometric function as follows

$$P^{(k,l)}_d(z) = \frac{(k+1)_d}{d!} {}_2F_1\left(-d, 1+k+l+d; k+1, \frac{1-z}{2}\right), \tag{1}$$

where $(k+1)_d$ is Pochhammer's symbol and ${}_2F_1$ is Gaussian or ordinary hypergeometric function defined for $|z| < 1$ by the power series

$${}_2F_1\left(a, b; c; z\right) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{n!(c)_n} z^n$$

Appell series F_1 defined by

$$F_1(a;b,b';c,x,y)=\sum_{n=0}^{\infty}\sum_{m=0}^{\infty}\frac{(a)_{m+n}(b)_n(b')_m}{n!m!(c)_{n+m}}x^n y^m,\quad (|x|<1,|y|<1)$$

is one of a natural two-variable extension of hypergeometric series ${}_2F_1$.
The following reduction formulas can be proved (see, for details [5], p. 238-239)

$$F_1(a;b,b';b+b';x,y) = (1-y)^{-a}{}_2F_1(a,b;b+b';(x-y)/(1-y)) \quad (2)$$
$$F_1(a;b,b';c;x,x) = {}_2F_1(a,b+b';c;x) \quad (3)$$

Comparing functions F_1 and $L_n^{2,2}$ it is easy to see

$$\frac{\Gamma(3+n)}{2}F_1(3+n;2,2;3;x,y)=L_n^{2,2}(x,y)$$

The following is the main theorem of this section.

Theorem 3.1 *The Bergman kernel for $D_n^{2,2}$ can be expressed in the following ways*

(i) $K_{D_n^{2,2}}((0,w_1,w_2),(0,\xi_1,\xi_2)) =$

$$\frac{\Gamma(3+n)}{2\pi^{n+2}}\sum_{i=0}^{n}\sum_{k=0}^{n-i}\binom{n}{i}\binom{n-1}{k}\frac{(2)_i(2)_k(w_1\overline{\xi}_1)^i(w_2\overline{\xi}_2)^k}{(3)_{i+k}(1-w_1\overline{\xi}_1)^{2+i}(1-w_2\overline{\xi}_2)^{2+k}}$$

(ii) *For every $m\in\mathbb{N}\cup\{0\}$, we have*

$$K_{D_{2m+1}^{2,2}}((0,w_1,w_2),(0,\xi_1,\xi_2)) = \frac{C_m(2+m)(2-\nu_1-\nu_2)P_m^{(\frac{3}{2},\frac{1}{2})}(\frac{x'+1}{1-x'})}{\pi^{2m+3}(1-\nu_1-\nu_2-\nu_1\nu_2)^{m+3}} - \frac{C_m(2m+1)P_m^{(\frac{3}{2},-\frac{1}{2})}(\frac{x'+1}{1-x'})}{\pi^{2m+3}(1-\nu_1-\nu_2-\nu_1\nu_2)^{m+2}},$$

where $\nu_1=w_1\overline{\xi}_1$, $\nu_2=w_2\overline{\xi}_2$, $x'=\left(\frac{\nu_1-\nu_2}{2-\nu_1-\nu_2}\right)^2$ and $C_m=\frac{\Gamma(4+2m)m!}{6(\frac{5}{2})_m}$.

(iii) *For every $m\in\mathbb{N}$, we have* $K_{D_{2m}^{2,2}}((0,w_1,w_2),(0,\xi_1,\xi_2)) =$

$$\Gamma(3+2m)m!\frac{2P_m^{(\frac{3}{2},-\frac{1}{2})}(\frac{x'+1}{1-x'})-(2-\nu_1-\nu_2)P_{m-1}^{(\frac{3}{2},\frac{1}{2})}(\frac{x'+1}{1-x'})}{\pi^{2m+2}6(\frac{5}{2})_{m-1}(1-\nu_1-\nu_2-\nu_1\nu_2)^{m+2}},$$

(iv) *If $w_1\overline{\xi}_1=w_2\overline{\xi}_2$ then*

$$K_{D_n^{2,2}}((0,w_1,w_2),(0,\xi_1,\xi_2))=\frac{\Gamma(3+n)(3+nw_1\overline{\xi}_1)}{6\pi^{n+2}(1-w_1\overline{\xi}_1)^{4+n}}$$

Proof. For the proof of (i), if we apply the recursion formula (see [8])

$$F_1(a+n;b,b';c;x,y) = \sum_{i=0}^{n}\sum_{k=0}^{n-i}\binom{n}{i}\binom{n-1}{k}\frac{(b)_i(b')_k}{(c)_{i+k}} \times x^i y^k F_1(a+i+k;b+i,b'+k;c+i+k;x,y),$$

then using formula

$$F_1(3+i+k;2+i,2+k;3+i+k;x,y) = \frac{1}{(1-x)^{2+i}(1-y)^{2+k}}$$

we obtain (i).
In order to prove the second statements we need the following well-known contiguous relation

$$cF_1(a;b,b';c;x,y) - (c-a)F_1(a;b,b';c+1;x,y) - aF_1(a+1;b,b';c+1;x,y) = 0$$

It follows that

$$F_1(4+2m;2,2;3;x,y) = -\frac{2m+1}{3}F_1\left(2m+4;2,2;4;x,y\right) + \frac{4+2m}{3}F_1\left(2m+5;2,2;4;x,y\right)$$

By (2), we have

$$F_1(4+2m;2,2;3;x,y) = -\frac{2m+1}{3(1-y)^{4+2m}}{}_2F_1\left(2m+4,2;4;\frac{x-y}{1-y}\right) + \frac{4+2m}{3(1-y)^{5+2m}}{}_2F_1\left(2m+5,2;4;\frac{x-y}{1-y}\right)$$

Then by (see, for details [5], p. 66)

$${}_2F_1(a,b;2b;z) = \left(1-\frac{z}{2}\right)^{-a}{}_2F_1\left(\frac{a}{2},\frac{a+1}{2};b+\frac{1}{2};[z/(2-z)]^2\right),$$

we have

$$F_1(4+2m;2,2;3;x,y) = -\frac{(2m+1)2^{4+2m}}{3(2-x-y)^{4+2m}}{}_2F_1\left(m+2,m+\frac{5}{2};\frac{5}{2};(x')^2\right) + \frac{(4+2m)2^{5+2m}}{3(2-x-y)^{5+2m}}{}_2F_1\left(m+\frac{5}{2},m+3;\frac{5}{2};(x')^2\right),$$

where $x' = \left(\frac{x-y}{2-x-y}\right)^2$. Next by (see, for details [5], p. 64)

$$\begin{aligned}{}_2F_1(a,b;c;z) &= (1-z)^{-a}\,{}_2F_1\left(a,c-b;c;z/(z-1)\right)\\ &= (1-z)^{-b}\,{}_2F_1\left(c-a,b;c;z/(z-1)\right),\end{aligned}$$

we can easily get $F_1(4+2m;2,2;3;x,y) =$

$$-\frac{2m+1}{3(1-x-y-xy)^{m+2}}{}_2F_1\left(m+2,-m;\frac{5}{2};\left(\frac{x'}{x'-1}\right)^2\right)$$
$$+\frac{(2+m)(2-x-y)}{3(1-x-y-xy)^{m+3}}{}_2F_1\left(m+\frac{5}{2},m+3;\frac{5}{2};\left(\frac{x'}{x'-1}\right)^2\right),$$

Finally, by (1) we obtain (ii). The formula (iii) can be obtained by the same method and we omit the details.
Now we will prove (iv). This is a formal exercise but we include it for completeness: From (3)

$$F_1(3+n;2,2;3;x,y) = {}_2F_1(3+n,4;3;x)$$

Now it is relatively easy to compute the following result

$${}_2F_1(3+n,4;3;x) = \frac{3+nx}{3(1-x)^{4+n}}$$

Thus we prove (iv).

4 Lu Qi-Keng problem

The explicit formula of the Bergman kernel function for the domains $D_n^{q,r}$ enables us to investigate whether the Bergman kernel has zeros in $D_n^{q,r} \times D_n^{q,r}$ or not. We will call this kind of problem a Lu Qi-Keng problem. If the Bergman kernel for a bounded domain does not have zeros, then the domain will be called a Lu Qi-Keng domain.

By Theorem 1.2 form [1] if $n = 1$, then $D_n^{q,r}$ is a Lu Qi-Keng domain for all positive real numbers q and r. Combining deflation identity (Proposition 2.3) and Proposition 4.4 from [1], if $n = 2$ and $q = r$, then $D_2^{r,r}$ is not a Lu Qi-Keng domain. Using the same method as in [1] we will prove that it is also true for $n = 3$.

Denote by

$$\begin{aligned}G(x,y) =& 3r^3(1-x)^3(1-y)^3 + 22r^2(1-x)^2(1-y)^2(1-xy)\\ &+ 24r(1-x)(1-y)\left(x(2x+1)y^2 + (x-8)xy + x + y + 2\right)\\ &+ 8(1-xy)\left(x^2y(4y+7) + x^2 + y^2 + xy(7y-38) + 7x + 7y + 4\right).\end{aligned}$$

Since

$$G(x,y) = \frac{r^3(1-x)^5(1-y)^5}{2}L_3^{r,r}(x,y)$$

then the Bergman kernel $K_{D_3^{r,r}}$ has zero inside $D_2^{r,r} \times D_2^{r,r}$ if polynomial $G(\epsilon x, \epsilon y)$ does not satisfy the stability property (see [1], for details) for some $0 < \epsilon < 1$. By following Šiljak and Stipanović [7] we consider polynomial

$$z^3G(e^{i\eta}, 1/z) = d(z) = d_3z^3 + d_2z^2 + d_1z + d_0,$$

where

$$
\begin{aligned}
d_0 &= 3r^3(-1+t)^3 - 22r^2(-1+t)^2t + 24rt(-1-t+2t^2) - 8t(1+7t+4t^2),\\
d_1 &= 8 - 9r^3(-1+t)^3 + 336t^2 - 56t^3 + 22r^2(-1+t)^2(1+2t)\\
&\quad - 24r(1-10t+8t^2+t^3),\\
d_2 &= 9r^3(-1+t)^3 - 22r^2(2-3t+t^3) - 8(-7+42t+t^3)\\
&\quad - 24r(1+8t-10t^2+t^3),\\
d_3 &= 22r^2(-1+t)^2 - 3r^3(-1+t)^3 - 24r(-2+t+t^2) + 8(4+7t+t^2)\\
&\text{and} \quad t = e^{i\eta}.
\end{aligned}
$$

With the polynomial $d(z)$ we associate the Schur-Cohn 3×3 matrix

$$
M(e^{i\eta}) = \begin{pmatrix} d_{11} & d_{12} & d_{13} \\ d_{21} & d_{22} & d_{23} \\ d_{31} & d_{32} & d_{33} \end{pmatrix},
$$

$d_{jk} = \sum_{l=1}^{j}(d_{m-j+l}\overline{d}_{m-k+l} - \overline{d}_{j-l}d_{k-l})$, where $1 \le j \le k$. The matrix $M(e^{i\eta})$ is defined when $j > k$ to become Hermitian. After some calculation (with the help of a computer program Maple or Mathematica), we have

$$
\det M(e^{i\eta}) = -\,130459631616r^3\sin^{12}(\eta/2)\left[\sum_{n=0}^{3} g_n(r)\cos(n\eta)\right],
$$

where

$$
\begin{aligned}
g_0(r) &= 26624 - 24672r^2 + 15724r^4 - 2430r^6\\
g_1(r) &= 12288 + 22496r^2 - 20822r^4 + 3645r^6\\
g_2(r) &= -512 + 2208r^2 + 5012r^4 - 1458r^6\\
g_3(r) &= r^2(-32 + 86r^2 + 243r^4)
\end{aligned}
$$

Since

$$
\sum_{n=0}^{3} g_n(r) = 38400
$$

Therefore it is easy to see that for every $r > 0$ there exist η such that $\det M(e^{i\eta}) < 0$. Hence there exist $1 > \epsilon > 0$, such that polynomial $G(\epsilon x, \epsilon y)$ does not satisfy the stability property (see [7]). As a consequence of above consideration, we have following corollary

Corollary 4.1 *For any $r > 0$, domains $D_3^{r,r}$ and $D_{1/3}^{r,r}$ are not Lu Qi-Keng.*

Now we will study Lu Qi-Keng problem for $D_n^{2,2}$ in the case when $n > 3$. By representation (iv) from Theorem 3.1

$$
K_{D_n^{2,2}}((0,w_1,w_1),(0,\xi_1,\xi_1)) = \frac{\Gamma(3+n)(3+nw_1\overline{\xi}_1)}{6\pi^{n+2}(1-w_1\overline{\xi}_1)^{4+n}}.
$$

Hence

$$K_{D_n^{2,2}}\left((0,i\sqrt{3/n},i\sqrt{3/n}),(0,-i\sqrt{3/n},-i\sqrt{3/n})\right)=0.$$

For brevity, we shall summarize these last statements by saying that

Proposition 4.1 *Domain $D_n^{2,2}$ is Lu Qi-Keng if and only if $n=1$.*

At the end of this section we would like to present following relations between zeros of the Bergman kernel for domains $D_n^{q,r}$ and $D_{1/n}^{q,r}$.

Proposition 4.2 *For any positive real numbers q and r*

(i) *If $K_{D_n^{q,r}}$ has zeros, then $K_{D_{1/n}^{q,r}}$ also has zeros,*

(ii) *If $D_{1/n}^{q,r}$ is Lu Qi-Keng domain, then $D_n^{q,r}$ is Lu Qi-Keng domain.*

Proof. Note that the zero set is a bi-holomorphic invariant object. Since any point $(z,w_1,w_2)\in D_n^{q,r}$ can be mapped equivalently onto the form $(0,\widetilde{w_1},\widetilde{w_2})$ by following automorphism of the $D_n^{q,r}$

$$D_n^{q,r}\ni(z,w_1,w_2)\mapsto\left(\Psi_a(z),\frac{(1-\|a\|^2)^{1/q}}{(1-\langle z,a\rangle)^{2/q}}w_1,\frac{(1-\|a\|^2)^{1/r}}{(1-\langle z,a\rangle)^{2/r}}w_2\right)\in\mathbb{C}^{n+2},$$

where

$$\Psi_a(z)=\frac{\left(\frac{\langle z,a\rangle}{1+\sqrt{1-\|a\|^2}}-1\right)a+z\sqrt{1-\|a\|^2}}{1-\langle z,a\rangle}.$$

Therefore, we need only consider the zeroes restricted to $\{0\}\times\mathbb{D}\times\mathbb{D}$, where $\mathbb{D}:=\{z\in\mathbb{C}:|z|<1\}$. The results now follows from Proposition 2.3.

5 Additional Results

Our purpose in this section is to consider domains Ω_n^r defined for every positive real number r by

$$\Omega_n^r:=\left\{(z,w)\in\mathbb{C}\times\mathbb{C}^n\colon |z|^2+|w_1|^r<1,\ldots,|z|^2+|w_n|^r<1\right\}$$

The reader can see that following proposition is completely analogous to the results presented earlier.

Proposition 5.1 *For $j=1,2,\ldots,n$ let $\beta_j\geq 0$. Then for $\alpha\geq 0$, we have*

$$\left\|z^\alpha w^\beta\right\|^2_{L^2(\Omega_n)}=\frac{\pi^{n+1}\Gamma\left(\frac{2}{r}(\sum_{j=1}^n\beta_j+n)+1\right)\Gamma(\alpha+1)}{\Gamma\left(\frac{2}{r}(\sum_{j=1}^n\beta_j+n)+\alpha+2\right)\prod_{j=1}^n(\beta_j+1)}.$$

Now we will give an explicit formula for the kernel $K_{\Omega_n^r}$ of Ω_n^r.

Theorem 5.1 *If $w, \zeta \in \mathbb{D}^n$, then the Bergman kernel for Ω_n^r is*

$$\pi^{n+1} K_{\Omega_n^r}((0,w),(0,\zeta)) = \prod_{k=1}^{n}(1-\nu_k)^{-2} + \sum_{k=1}^{n} \frac{2(1+\nu_k)}{r(1-\nu_1)^2\cdots(1-\nu_n)^2(1-\nu_k)},$$

where $\nu_k = w_k\overline{\zeta}_k$ for $k = 1, \ldots, n$. Moreover if $\nu_1 = \nu_2 = \cdots = \nu_n$, then

$$\pi^{n+1} K_{\Omega_n^r}((0,w),(0,\zeta)) = \frac{(2n-r)\nu_1 + 2n + r}{r(1-\nu_1)^{2n+1}}.$$

Proof. As a consequence of Proposition 5.1, we have

$$\pi^{n+1} K_{\Omega_n^r}((0,w),(0,\zeta)) = \sum_{\beta_1,\ldots,\beta_n \geq 0}^{\infty} \frac{\Gamma\left(\frac{2}{r}(\sum_{j=1}^n \beta_j + n) + 2\right) \prod_{j=1}^n (\beta_j+1)}{\Gamma\left(\frac{2}{r}(\sum_{j=1}^n \beta_j + n) + 1\right)} \nu^\beta,$$

where $\nu^\beta = \nu_1^{\beta_1} \cdot \cdots \cdot \nu_n^{\beta_n}$. Then, using the fact that $a\Gamma(a) = \Gamma(a+1)$,

$$\pi^{n+1} K_{\Omega_n^r}((0,w),(0,\zeta)) = \sum_{\beta_1,\ldots,\beta_n \geq 0}^{\infty} \frac{2}{r}\left(\beta_1 + \cdots + \beta_n + n + \frac{r}{2}\right) \prod_{j=1}^{n} (\beta_j+1)\nu^\beta.$$

Hence, the kernel is

$$\frac{2}{r\pi^{n+1}} \sum_{k=0}^{n} \sum_{\beta_1,\ldots,\beta_n \geq 0}^{\infty} (\beta_k+1) \prod_{j=1}^{n} (\beta_j+1)\nu^\beta,$$

where $\beta_0 = r/2 - 1$. After some calculation using formulas

$$\sum_{m=0}^{\infty} (m+1)x^m = (1-x)^{-2} \quad \text{and} \quad \sum_{m=0}^{\infty} (m+1)^2 x^m = \frac{1+x}{(1-x)^3},$$

we obtain desired formula.

5.1 Zeros of Bergman kernels on Ω_n^r

In this section, we will prove that the Bergman kernel function of Ω_n^r for any natural number n and positive real number r is zero-free.

Proposition 5.2 *For any positive integer n, the domain Ω_n^r is a Lu Qi-Keng domain.*

Proof. It is obvious that the Bergman kernel for Ω_1^r has no zeros. The Bergman kernel function of Ω_n^r has zeros iff

$$\frac{r}{2} + \sum_{k=1}^{n} \frac{1+\nu_k}{1-\nu_k} = n + \frac{r}{2} + \sum_{k=1}^{n} \frac{2\nu_k}{1-\nu_k} = 0,$$

for some $\nu_1, \ldots, \nu_n$ such that $|\nu_k| < 1$, for each $k = 1, \ldots, n$. Since $|\nu_k| < 1$, then

$$\Re\left(\frac{2\nu_k}{1-\nu_k}\right) > -1,$$

where $\Re(\xi)$ is the real part of complex number ξ. Thus

$$\Re\left(n + \frac{r}{2} + \sum_{k=1}^{n} \frac{2\nu_k}{1-\nu_k}\right) > 0.$$

Hence, we see that

$$\frac{r}{2} + \sum_{k=1}^{n} \frac{1+\nu_k}{1-\nu_k} \neq 0,$$

if $(\nu_1, \ldots, \nu_n) \in \mathbb{D}^n$. The proof is therefore complete.

References

[1] T. Beberok, Lu Qi-Keng's Problem for Intersection of Two Complex Ellipsoids, *Complex Anal. Oper. Theory.* DOI: 10.1007/s11785-015-0505-4 (2015).

[2] S. Bergman, Zur Theorie von pseudokonformen Abbildungen, *Mat. Sb. (N.S.)* 1(43)(1) (1936) 79–96.

[3] H.P. Boas, S. Fu and E.J. Straube, The Bergman kernel function: explicit formulas and zeroes, *Proc. Amer. Math. Soc.* **127**(3) (1999) 805–811.

[4] J.P. D'Angelo, An explicit computation of the Bergman kernel function, *J. Geom. Anal.* **4** (1994) 23–34.

[5] A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions*, (vol. 1, McGraw-Hill, New York, 1953).

[6] S. G. Krantz, *Geometric analysis of the Bergman Kernel and Metric*, (Springer, New York, 2013).

[7] D. D. Šiljak and D. M. Stipanović, Stability of Interval Two-Variable Polynomials and Quasipolynomials via Positivity, *Lecture Notes in Control and Information Sciences.* **312** (2005) 165–177.

[8] X. Wang, Recursion formulas for Appell functions, *Integral Transforms Spec. Funct.* **23**(6) (2012) 421–433.

Tomasz Beberok
Department of Applied Mathematics
University of Agriculture in Krakow
ul. Balicka 253c, 30-198 Krakow, Poland
email: tbeberok@ar.krakow.pl